

Ali-Reza Ketabi, Sandra Ketabi, Hans-Christoph Lauer, Andree Piwowarczyk, Martin Brenner

# Are apical lesions visible more often in CBCT than in panoramic radiographs?

## Abstract

**Introduction:** This study compares the accuracy of panoramic radiography and CBCT in detecting and diagnosing mandibular apical lesions, analysing the effect of the thickness of the cortical bone on the radiological visibility of apical lesions.

**Methods:** Digital images of 343 patients who had CBCT scans and panoramic radiographs were inspected for mandibular apical lesions. The dimensions of the lesion and the thickness of the cortical bone in the affected region were measured. Statistical analyses were made using statistical software (IBM SPSS 25, IBM, Armonk, NY, USA).

**Results:** CBCT detected apical lesions around 115 teeth; 77 of these were also visible on a panoramic radiography. The differences between the two methods were statistically significant in the premolar and molar regions (McNemar test;  $p < 0.001$ ) but not in the anterior region ( $p = 0.063$ ). For the size of the lesions no significant differences were found (Wilcoxon test; anterior,  $p = 0.60$ ; premolar,  $p = 0.90$ ; molar,  $p = 0.61$ ). In the Mann-Whitney U test, buccal and palatal cortical thickness did not significantly influence the visibility of lesions in CBCT and panoramic measurements (buccal,  $p = 0.93$ ; palatal,  $p = 0.15$ ).

**Conclusion:** Apical lesions seem to be much more readily visible on CBCT scans. The thickness of the cortical bone and the size of the lesions do not significantly influence the representation of apical lesions.

**Keywords:** apical lesions; CBCT; cone-beam computed tomography; cortical bone thickness; panoramic radiography

Department of Prosthodontics, School of Dentistry, Faculty of Health, Witten/Herdecke University, Alfred-Herrhausen-Straße 45, 58455 Witten, Germany: Dr Ali-Reza Ketabi, Prof Dr Andree Piwowarczyk

Private Dental Office of Dr Ali-Reza Ketabi, Epplestraße 29 a, 70597 Stuttgart, Germany: Dr Ali-Reza Ketabi, Dr Sandra Ketabi

Department of Prosthodontics, Center for Dentistry and Oral Medicine (Carolinum), Goethe University, Theodor-Stern-Kai 7, 60596 Frankfurt, Germany: Prof Dr Hans-Christoph Lauer

Private Dental Office of Dr Martin Brenner, Zahnärzte am Schloss, Bismarckstr. 15, 72574 Bad Urach, Germany: Dr Martin Brenner

**Citation:** Ketabi A-R, Ketabi S, Lauer H-C, Piwowarczyk P, Brenner M: Are apical lesions visible more often in CBCT than in panoramic radiographs? *Dtsch Zahnärztl Z Int* 2022; 4: 164–170

**Peer reviewed article:** submitted: 09.11.2021, revised version accepted: 21.04.2022

**DOI.org/10.53180/dzz-int.2022.0020**

## Introduction

Apical bony lesions, usually detected as a radiolucency around the root on radiographs, are a common sequela of endodontic infections [2]. Selecting the appropriate therapeutic approach requires an accurate diagnosis [7]. The diagnosis, assessment and treatment of an endodontic condition is generally accomplished by periapical radiography [2, 8] or panoramic radiography [1]. The current gold standard for detection of periapical lesions is routinely periapical radiography [2, 8]. Practitioners often prefer digital imaging because it requires no time-consuming processing [18, 21, 24] and the results are more accurate. Digital imaging is more sensitive in detecting apical lesion than conventional periapical radiography [9, 10]. However, there have been frequent reports that bony aspects of apical bony lesions were not always detectable on periapical radiographs [31]. If these lesions involve cortical bone (buccal or lingual), this will typically result in reduced radiodensity in some areas, whereas lesions in cancellous bone enclosed by dense cortical bone will frequently remain undetected [7, 12, 14, 28].

The most important factor that determines whether apical bony lesions are visible on intraoral radiographs is the presence or absence of cortical perforation. Van Assche et al. found that approximately 90% of these lesions were detected if the cortex had been perforated, but the examiners detected only 10% of intrabony lesions [27]. A correct diagnosis is also made more difficult by variations in apical morphology, x-ray angulations, bone density and radiographic contrast [11].

Three-dimensional (3D) imaging is an alternative to two-dimensional (2D) panoramic and periapical radiography in detecting apical lesions. The 3D technology most frequently used is cone-beam computed tomography (CBCT). Recent studies have looked at the diagnostic value of CBCT scans in relation to apical lesions [2, 3, 14, 20, 26, 29]. Paula-Silva et al., comparing periapical radiographs with CBCT and histological findings, found that the apical lesions were not detected on periapi-



Figures 1–3: Ali-Reza Ketabi

**Figure 1** By panoramic radiography (a), apical lesions are not visible on tooth 45 and barely visible on the mesial root of tooth 47, whereas they are clearly perceptible on teeth 45 (b) and 47 (c) by CBCT taken 5 days later.

cal radiographs in 22% of cases, while CBCT only failed to detect them in 9% of cases [6]. This was confirmed by studies reporting significantly more accurate diagnoses for CBCT than had been the case for periapical radiographs [2, 7, 14]. It should be noted, however, that most of the studies were either ex vivo or animal studies [2, 6, 14], had small study populations [3, 26] or did not directly compare 2D images and 3D images for specific patients [7, 20, 29]. A systematic review with meta-analysis by Leonardi Dutra et al. yielded no data for comparing panoramic radiography and CBCT [17].


The aim of the present study, therefore, was to compare the visibility of mandibular apical lesions in 2D (panoramic radiography) and 3D (CBCT) radiographs of the same patient and determine whether cortical bone thickness influenced the detectability of apical lesions in a study population of adequate size. Furthermore, it was the authors' intention to measure the size and number of the lesions.

## Material and methods


Ethical approval was secured from the Medical Council of Baden-Württemberg, Germany (Register No.

Indication	Gender		
	Female	Male	Total
Endodontics	13 7.5%	10 5.9%	23 6.7%
Periodontology	15 8.7%	8 4.7%	23 6.7%
Implantology	118 68.2%	118 69.8%	236 69.0%
Miscellaneous	103 59.5%	86 50.9%	189 55.3%

**Table 1** Distribution of justifying indications for CBCT



**Figure 2** The measurements of the height AP on the PR and the CBCT.



**Figure 3** The measurements of the width AP on the PR and the CBCT.

F-2014–006-z). The present study was conducted in accordance with the ethical standards of the 1964 Declaration of Helsinki [32]. Imaging records for over 1,000 patients (including panoramic radiography and CBCT) were selected from a data repository at a dental radiological centre. The original examinations had taken place between February of 2010 and January of 2017. Endodontic, periodontal and implant-related diagnoses were included, in addition to some miscellaneous indications (such as a search for cysts, impacted teeth or bone fractures). All patients that were included had had at least one panoramic radiograph and one CBCT scan taken within the previous three months. Patients with an interval between examinations of more than three months were excluded to avoid any distortion potentially caused by changes in the anatomy (such as the development of apical lesions). Data were anonymised prior to analysis.

Digital panoramic radiographs were obtained using an Orthophos D3297 (Sirona, Bensheim, Germany) with 60 kVp and a size of the focal spot of 0.5 mm. Panoramic radiographs were calibrated by reference to objects with a known length (such as implants, metal objects, root canal-treated restored teeth) to eliminate any errors that might have been caused by magnification. Cases without reference objects were excluded. CBCT scans were acquired with a Gendex GXCB-500 (KaVo, Hamburg, Germany) with an 8 × 8 cm field of view (FoV). The exposure was set at 120 kVp/5 mA and 6.0 seconds; the voxel size was set at 0.2 mm. All images were obtained by a dental radiologist. The panoramic images were evaluated with imaging software (DBSWIN v. 5.1.1, Dürr Dental, Bietigheim-Bissingen, Germany). CBCT images were examined using i-CAT Vision software (Imaging Sciences, Hatfield, PA, USA). One examiner with the necessary specialist CBCT competence visually analysed the digital images on a validated computer monitor (Eizo Flex Scan S2000, Eizo Corp., Hakusan, Ishikawa, Japan) in a darkroom after being

specially trained by a dental radiologist.

After the training, inter-examiner and intra-examiner reliability values were calculated for 20 samples. The measurements proceeded for no more than 6 hours per day, with 30-minute periods of rest every 2 hours. All records were reviewed for errors and corrected by a dental radiologist if necessary. Only teeth visible on both the panoramic radiograph and CBCT were included.

The presence or absence of apical lesions was determined for each tooth (Fig. 1), with lesion size measured in vertical (Fig. 2) and horizontal (Fig. 3) direction. For CBCT images, the buccopalatal extension of any lesions and the cortical bone thickness (buccally and palatally) in the area of the lesions, if appropriate, were measured.

Data were transferred to individual data acquisition forms for analysis by statistical software (SPSS 25) by a professional medical-statistics provider (medistat, Kronshagen, Germany). The frequency of apical lesions was recorded separately for anteriors/canines, premolars and molars.

The working hypothesis was: "There is no difference in the visibility of apical lesions on panoramic radiographs and CBCT scans."

The CBCT and PR measurements were tested by the Wilcoxon Test for Pair Differences for variations. Two independent samples were compared by means of the Mann-Whitney U test. To examine correlations between quantitative, not normally distributed parameters, a rank correlation analysis was carried out as per Spearman.

The McNemar test and the chi-square test were used for a statistical testing of the working hypothesis (differences between imaging techniques regarding the visibility of periapical lesions). A p-value of  $\leq 0.05$  indicates the presence of a significant difference.

**Results**

A total of 343 patients for which mandibular panoramic radiographs and CBCT scans – taken within 3 months – were available were in-

Tooth group			CBCT		Total
			Negative	Positive	
Anterior teeth	Panoramic	Negative	1,043 98.2%	5 0.5%	1,048 98.7%
		Positive	0 0.0%	14 1.3%	14 1.3%
		Total	1,043 98.2%	19 1.8%	1,062 100%
Premolars	Panoramic	Negative	575 92.7%	18 2.9%	593 95.6%
		Positive	0 0.0%	27 4.4%	27 4.4%
		Total	575 92.7%	45 7.3%	593 100%
Molars	Panoramic	Negative	578 91.9%	15 2.4%	593 94.3%
		Positive	0 0.0%	36 5.7%	36 5.7%
		Total	578 91.9%	51 8.1%	629 100%
Total	Panoramic	Negative	2,196 95.0%	38 1.6%	2,234 96.7%
		Positive	0 0.0%	77 3.3%	77 3.3%
		Total	2,196 95.0%	115 5.0%	2,311 100.0%

Computed tomography (CBCT) versus panoramic radiographs in mandible.  
**Table 2** Cross-classified comparison of the visibility of apical bony lesions by cone-beam

cluded in the study. Genders were represented approximately equally (174 female/169 male). The patients' age at baseline was  $59.0 \pm 14.5$  (mean  $\pm$  SD; range 19–92) years. The indications for CBCT scans are shown in Table 1; in some cases, there were multiple indications. The inter-examiner and intra-examiner reliabilities were rated as very high (Cohen's kappa 1.0; 95% CI for kappa [0.92; 1.00]).

The prevalence of apical lesions as identified by panoramic radiographs and CBCT is presented as a cross-classified table (Table 2). 67.0% of the lesions visible by CBCT were also visible on a panoramic radiography. The visibility differences were significant in the premolar and molar regions (McNemar test;  $p < 0.001$ )

but not in the anterior region ( $p = 0.063$ ).

The size of the apical lesions was also determined (Table 3). Some teeth were affected by multiple lesions. Hence, the total number of lesions included in the measurements was 123 (115 teeth). The CBCT measurements were: width  $3.38 \pm 3.11$  mm (mean  $\pm$  SD; range 0.90–33.30 mm); height  $1.95 \pm 1.47$  mm; (0.00–11.60 mm); depth  $3.33 \pm 1.48$  mm (1.20–9.00 mm). The panoramic measurements were: width  $3.63 \pm 3.04$  mm (0.80–29.4 mm); height  $1.50 \pm 1.22$  mm (0.00–10.30 mm). The Wilcoxon test showed no significant differences between the two imaging technologies (anteriors/canines  $p = 0.60$ ; premolars  $p = 0.90$ ; molars  $p = 0.61$ ).

		N	Mean	SD	Min.	Max.	25 <sup>th</sup>	50 <sup>th</sup> (Median)	75 <sup>th</sup>
<b>Anterior teeth</b>	CBCT: width	20	3.06	1.92	1.20	9.80	2.10	2.40	3.60
	CBCT: height	20	2.06	2.35	0.60	11.60	1.13	1.30	2.08
	CBCT: depth	20	3.38	1.66	1.80	9.00	2.40	2.85	3.85
	CBCT: buccal cortical bone thickness	20	1.39	0.51	0.00	2.40	1.20	1.50	1.50
	CBCT: palatal cortical bone thickness	20	1.89	0.58	0.90	3.10	1.50	1.80	2.18
	PANO: width	17	3.56	1.90	1.10	9.40	2.10	3.50	4.15
	PANO: height	17	1.89	2.31	0.50	10.30	0.60	1.20	2.25
	<b>Premolars</b>	CBCT: width	46	2.92	1.17	0.90	6.30	2.10	2.70
CBCT: height		46	1.68	1.01	0.00	5.10	1.20	1.50	2.10
CBCT: depth		46	2.70	0.96	1.20	5.40	2.08	2.40	3.33
CBCT: buccal cortical bone thickness		46	1.85	0.57	0.00	3.00	1.50	1.80	2.18
CBCT: palatal cortical bone thickness		46	2.00	0.52	1.20	3.40	1.50	1.80	2.13
PANO: width		37	3.34	1.34	1.40	6.00	2.15	3.30	4.45
PANO: height		36	1.34	0.88	0.00	3.70	0.60	1.05	2.00
<b>Molars</b>		CBCT: width	57	3.90	4.32	1.10	33.30	2.10	3.00
	CBCT: height	57	2.14	1.39	0.70	7.60	1.20	1.80	2.70
	CBCT: depth	57	3.82	1.60	1.50	8.80	2.65	3.30	4.70
	CBCT: buccal cortical bone thickness	57	2.70	0.89	0.00	4.50	2.15	2.70	3.35
	CBCT: palatal cortical bone thickness	57	2.06	0.71	0.00	3.70	1.50	2.10	2.50
	PANO: width	53	3.86	4.05	0.80	29.40	2.00	3.20	4.20
	PANO: height	53	1.49	0.87	0.40	3.80	0.90	1.20	1.80

**Table 3** Measurements of apical lesion parameters and cortical bone thickness in the mandible

Mean cortical thickness, as measured by CBCT, was  $2.17 \pm 0.71$  mm (mean  $\pm$  SD; range 0.00–4.50 mm) buccally and  $2.01 \pm 0.62$  mm (0.00–3.70 mm) palatally. The results for the thickness of compact bone near apical lesions are given in Table 3. In the Mann-Whitney U test, buccal and palatal cortical thickness did not significantly influence the visibility of lesions in CBCT and panoramic measurements (buccal  $p = 0.93$ ; palatal  $p = 0.15$ ).

The cortical bone was significantly thicker both on the buccal and on the palatal aspect (Mann-Whitney U test;  $p < 0.01$ ) for molars on which apical lesions were visible on both

CBCT and panoramic radiographs. The visibility of the apical lesions showed no correlation with bone thickness in the anterior (buccal  $p = 0.65$ ; palatal  $p = 0.59$ ) and premolar regions (buccal  $p = 0.68$ ; palatal  $p = 0.11$ ).

### Discussion

The results show that apical lesions were 1.5 times more visible on CBCT scans than on panoramic radiographs. Furthermore, the diameters of the apical lesions were almost equivalent for both imaging methods.

This result agrees with a study by Nardi et al., who investigated the ac-

curacy of panoramic radiography in detecting apical lesions on asymptomatic root canal-treated teeth and found higher positive predictive values for lesions located in the mandible in canine/premolar and molar areas [19]. Furthermore, Nardi et al. found negative predictive values for panoramic radiography if the lesions were smaller than 4.6 mm and did not affect the cortical bone [19]. In the present study, discrepancies were highly evident. A possible explanation in the anterior region is that the anterior imaging quality is often poor in the latter due to a possible superimposition of the spine or blurring caused by a position out of

focus. Hence, this value is significantly higher than the visibility of anterior maxillary lesions, of which only 13.6% were detected via panoramic radiography. This discrepancy is caused by the roots in the mandible being more orthograde to the x-ray beam. Furthermore, disturbances by superimposition of anatomical structures are less common in the mandible [19]. Not once was a lesion found on a panoramic radiograph if not on the corresponding CBCT. However, not least because of radiation protection requirements in accordance with the ALARA/ALADA principle [13] (“As low as reasonably achievable/as low as diagnostically acceptable”), panoramic radiography is not the diagnostic method of choice for apical lesions [19].

The present study is of limited validity because histological material could not be acquired for ethical reasons. False-negative/false-positive results may have been present. However, histopathologic studies reached comparable results and showed high diagnostic accuracy for CBCT and apical lesions compared with panoramic radiography [2]. Other human clinical studies used CBCT as reference, [4, 5, 7, 22, 23] with possibly biased results [25, 30]. Recently, Kruse et al. found that the diagnostic accuracy of CBCT depends on the endodontic status of the tooth. The diagnostic accuracy of CBCT was high for non-treated roots, whereas the diagnosis of apical lesions on root canal-treated roots was less accurate [16]. This aspect had not been known when the present study was implemented and was therefore not considered.

For this study the researcher was briefed by an expert in the field of dental radiology prior to commencement of the study. To verify the reliability of radiographic measurements and evaluations, multiple ratings were carried out of 20 randomly selected patients. Since a selected patient population was included here, a possible risk of spectrum bias cannot be ruled out. However, the high intra- and interrater reliability (Cohen’s kappa 1.0) of the study also indicates the reliability of the results.

A systematic review and meta-analysis by Leonardi Dutra et al., on

the other hand, found high levels of accuracy for CBCT (0.96) compared to the already good results obtained by periapical radiography (0.72–0.73) in detecting apical lesions. The present results combined with current published research allow us to conclude that CBCT is more accurate in detecting apical lesions than panoramic radiography [17].

The thickness of the cortical bone and its influence on the visibility of lesions on CBCT and panoramic radiography has not been investigated previously for the mandible. Our results do not show any significant correlation between the thickness of the cortical bone and the visibility of lesions on CBCT and panoramic radiography. The cortical bone in the molar area was significantly thicker in cases where apical lesions were detected, as compared to cases where no such lesions were visible. It is conceivable that the thicker cortical bone compensates for the loss of cancellous bone in the maxillary molar region with its typically lower bone density. However, this would need to be verified in further studies.

There was no correlation between cortical bone perforation and lesion visibility in the mandible, contradicting previous results [27].

The results show that panoramic radiographs are not a reliable diagnostic tool for detecting apical lesions. Small apical lesions seem to be much better visible on CBCT scans. Cortical thickness does not seem to influence the visibility of apical lesions on panoramic radiographs or CBCT scans.

### Authorship declaration

Dr Ali-Reza Ketabi had the idea for the study and collected the data. Dr Sandra Ketabi evaluated the radiographs and was responsible for the data analysis. Prof Dr Hans-Christoph Lauer substantively revised the manuscript. Prof Dr Andree Piowarczyk substantively revised the manuscript. Dr Martin Brenner wrote the manuscript and provided support in the conception of the study. All authors are in agreement with the manuscript.

### Conflict of interest

The authors declare that there is no conflict of interest as defined by the guidelines of the International Committee of Medical Journal Editors.

### Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

### References

1. Bender IB: Factors influencing the radiographic appearance of bony lesions. *J Endod.* 1982; 8(4): 161–170 doi: 10.1016/S0099-2399(82)80212-4. PMID: 6951916.
2. Campello AF, Goncalves LS, Guedes FR, Marques FV: Cone-beam computed tomography versus digital periapical radiography in the detection of artificially created periapical lesions: a pilot study of the diagnostic accuracy of endodontists using both techniques. *Imaging Sci Dent* 2017; 47: 25–31.
3. Chanani A, Adhikari HD: Reliability of cone beam computed tomography as a biopsy-independent tool in differential diagnosis of periapical cysts and granulomas: an in vivo Study. *J Conserv Dent* 2017; 20: 326–331.
4. Davies A, Mannocci F, Mitchell P, Andiappan M, Patel S: The detection of periapical pathoses in root filled teeth using single and parallax periapical radiographs versus cone beam computed tomography – a clinical study. *Int Endod J* 2015; 48: 582–592.
5. Davies A, Patel S, Foschi F, Andiappan M, Mitchell PJ, Mannocci F: The detection of periapical pathoses using digital periapical radiography and cone beam computed tomography in endodontically retreated teeth – part 2: a 1 year post-treatment follow-up. *Int Endod J* 2016; 49: 623–635.
6. de Paula-Silva FW, Wu MK, Leonardo MR, da Silva LA, Wesselink PR: Accuracy of periapical radiography and cone-beam computed tomography scans in diagnosing apical periodontitis using histopath-

ological findings as a gold standard. *J Endod* 2009; 35: 1009–1012.

7. Estrela C, Bueno MR, Leles CR, Azevedo B, Azevedo JR. Accuracy of cone beam computed tomography and panoramic and periapical radiography for detection of apical periodontitis. *J Endod* 2008; 34: 273–279.
8. European Society of Endodontology: Quality guidelines for endodontic treatment: consensus report of the European Society of Endodontology. *Int Endod J* 2006; 39: 921–930.
9. Folk RB, Thorpe JR, McClanahan SB, Johnson JD, Strother JM: Comparison of two different direct digital radiography systems for the ability to detect artificially prepared periapical lesions. *J Endod* 2005; 31: 304–306.
10. Hadley DL, Replogle KJ, Kirkam JC, Best AM: A comparison of five radiographic systems to D-speed film in the detection of artificial bone lesions. *J Endod* 2008; 34: 1111–1114.
11. Halse A, Molven O, Fristad I: Diagnosing periapical lesions--disagreement and borderline cases. *Int Endod J* 2002; 35: 703–709.
12. Huuromonen S, Ørstavik D. Radiological aspects of apical periodontitis. *Endod Topics* 2002; 1: 3–25.
13. Jaju PP, Jaju SP: Cone-beam computed tomography: time to move from ALARA to ALADA. *Imaging Sci Dent* 2015; 45: 263–265.
14. Kanagasingam S, Lim CX, Yong CP, Mannocci F, Patel S: Diagnostic accuracy of periapical radiography and cone beam computed tomography in detecting apical periodontitis using histopathological findings as a reference standard. *Int Endod J* 2017; 50: 417–426.
15. Kruse C, Spin-Neto R, Reibel J, Wenzel A, Kirkevang LL: Diagnostic validity of periapical radiography and CBCT for assessing periapical lesions that persist after endodontic surgery. *Dentomaxillofac Radiol* 2017; 46: 20170210.
16. Kruse C, Spin-Neto R, Evar Kraft DC, Vaeth M, Kirkevang LL: Diagnostic accuracy of cone beam computed tomography used for assessment of apical periodontitis: an ex vivo histopathological study on human cadavers. *Int Endod J* 2019; 52: 439–450.
17. Leonardi Dutra K, Haas L, Porporatti AL, Flores-Mir C, Nascimento Santos J, Mezzomo LA et al.: Diagnostic accuracy of cone-beam computed tomography and conventional radiography on apical periodontitis: a systematic review and meta-analysis. *J Endod* 2016; 42: 356–364.
18. Nair MK, Nair UP: Digital and advanced imaging in endodontics: a review. *J Endod* 2007; 33: 1–6.
19. Nardi C, Calistri L, Grazzini G, Desideri I, Lorini C, Occhipinti M, et al: Is panoramic radiography an accurate imaging technique for the detection of endodontically treated asymptomatic apical periodontitis? *J Endod* 2018; 44: 1500–1508.
20. Nascimento EH L, Oenning ACC, Freire BB, Gaeta-Araujo H, Haiter-Neto F, Freitas DQ: Comparison of panoramic radiography and cone beam CT in the assessment of juxta-apical radiolucency. *Dentomaxillofac Radiol* 2018; 47: 20170198.
21. Patel S, Dawood A, Whaites E, Pitt Ford T: New dimensions in endodontic imaging: part 1. Conventional and alternative radiographic systems. *Int Endod J* 2009; 42: 447–462.
22. Patel S, Wilson R, Dawood A, Foschi F, Mannocci F: The detection of periapical pathology using digital periapical radiography and cone beam computed tomography – Part 2: a 1-year post-treatment follow-up. *Int Endod J* 2012; 45: 711–723.
23. Patel S, Wilson R, Dawood A, Mannocci F: The detection of periapical pathology using periapical radiography and cone beam computed tomography – Part 1: pre-operative status. *Int Endod J* 2012; 45: 702–710.
24. Peters CI, Peters OA: Cone beam computed tomography and panoramic and periapical radiography for detection of apical periodontitis. *Endod Topics* 2012; 34: 57–75.
25. Pope O, Sathorn C, Parashos P: A comparative investigation of cone-beam computed tomography and periapical radiography in the diagnosis of a healthy periapex. *J Endod* 2014; 40: 360–365.
26. Torabinejad M, Rice DD, Maktabi O, Oyoyo U, Abramovitch K: Prevalence and size of periapical radiolucencies using cone-beam computed tomography in teeth without apparent intraoral radiographic lesions: a new periapical index with a clinical recommendation. *J Endod* 2018; 44: 389–394.
27. Van Assche N, Jacobs R, Coucke W, van Steenberghe D, Quirynen M: Radiographic detection of artificial intra-bony defects in the edentulous area. *Clin Oral Implants Res* 2009; 20: 273–279.
28. van der Stelt PF: Experimentally produced bone lesions. *Oral Surg Oral Med Oral Pathol* 1985; 59: 306–312.
29. Van der Veken D, Curvers F, Fieuws S, Lambrechts P: Prevalence of apical periodontitis and root filled teeth in a Belgian subpopulation found on CBCT images. *Int Endod J* 2017; 50: 317–329.
30. Walter SD, Macaskill P, Lord SJ, Irwig L: Effect of dependent errors in the assessment of diagnostic or screening test accuracy when the reference standard is imperfect. *Stat Med* 2012; 31(11–12): 1129–1138.
31. White SC, Atchison KA, Hewlett ER, Flack VF: Efficacy of FDA guidelines for prescribing radiographs to detect dental and intraosseous conditions. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 1995; 80: 108–114.
32. World Medical Association: World Medical Association Declaration of Helsinki: ethical principles for medical research involving human subjects. *JAMA* 2013; 310: 2191–2194.



Photo: A.-R. Ketabi

#### Corresponding author

**DR ALI-REZA KETABI**

Dentist – Department of Prosthodontics, School of Dentistry, Faculty of Health, Witten/Herdecke University, 58455 Witten, Germany and Private Dental Office of Dr Ali-Reza Ketabi, Epplestraße 29 a, 70597 Stuttgart, Germany

<https://orcid.org/0000-0001-5752-2529>

E-mail: [ali-reza.ketabi@uni-wh.de](mailto:ali-reza.ketabi@uni-wh.de), [alirezaketabi@yahoo.de](mailto:alirezaketabi@yahoo.de)

Phone: +49 711 6339697-0

Fax: +49 711 6339697-7