

Matthias G. Hautmann

# Position paper: Does radiotherapy play a role in activated osteoarthritis of the temporomandibular joint?

**Topic: Could radiotherapy be a treatment option for symptomatic inflammatory and/or degenerative arthropathy of the temporomandibular joint?**



## Background

In addition to radiotherapy for malignant tumors, radiotherapy for non-malignant diseases also plays an important role in daily clinical practice and it is applied successfully [17, 24]. Although radiotherapy is used much more frequently to treat malignant diseases, several national patterns of care studies have shown the relevance of radiotherapy for treating non-malignant diseases. The last large national patterns of care study revealed that more than 37,000 patients per year are irradiated for non-malignant diseases. In this case, only a small proportion are benign tumors. The majority of cases consist of degenerative and degenerative-inflammatory diseases, followed by functional diseases and hyperproliferative diseases. Degenerative and degenerative-inflammatory diseases alone account for more than 23,000 cases per year in Germany [25]. Consequently, the interesting question arises of whether patients with degenerative or degenerative-inflammatory diseases of the TMJ can also benefit from radiotherapy.

Among the degenerative and degenerative-inflammatory diseases, plantar fasciitis (with or without accompanying heel spur), enthesiopathy, tendinitis, epicondylitis, and subacromial syndrome play an important role [17, 24, 25]. In a large percentage, many of these diseases can be cured or at least significantly improved by means of radiotherapy. For example, in plantar fasciitis, radiotherapy is a standard therapy. The effect has been demonstrated in numerous prospective, randomized studies [18, 24].

Radiotherapy for the treatment of osteoarthritis has been an established procedure for many years. The beginnings of radiotherapy in the treatment of arthritic changes dates back to the 1920s [11, 15, 16, 28]. However, there is little data on radiotherapy for osteoarthritis or activated osteoarthritis of the TMJ. Most existing data relates to knee joint arthritis, hip arthritis and finger polyarthritits. In many heterogeneous collectives, TMJ osteoarthritis is lost in the “other” arthritides. Only one older study reports TMJ osteoarthritis separately [2, 4, 11].

The current S2k-Guideline of the DEGRO (German Society for Radiation

Oncology) provides a “should” recommendation for gonarthritits and a “can” recommendation for other arthritides. Thus, according to the guideline’s recommendations, there is also the possibility to employ irradiation in the treatment of osteoarthritis or degenerative and/or inflammatory arthropathies of the TMJ [16, 17, 20]. The author’s research group has been successfully treating patients with TMJ osteoarthritis using radiotherapy for several years. Most of these patients are included in a national multicenter observational study.

## Principles of radiobiology

Many preclinical studies have shown that low-dose radiotherapy has an inflammation-modulating or anti-inflammatory effect via different mechanisms. For example, a reduction of chemotactic cytokines (e.g., CCL20) has been shown. Thus, fewer inflammatory cells are attracted [17, 22].

One of the most important stages in the inflammatory cascade is the adhesion of monocytes and granulocytes to the endothelium and the migration of these immune cells into the inflamed tissue. One mode of ac-




Fig. 1: M. C. Hautmann

**Figure 1** Radiotherapy plan for a patient who has been irradiated unilaterally at the TMJ. The figures on the left and above show the dose distribution in various 3D cross sections: two transversal, one sagittal and one coronal. The target volume is outlined in red. The figure on the bottom right shows a dose-volume histogram which is used to estimate the dose distribution in the target volume and the dose exposure of the organs at risk.

tion of low-dose radiotherapy is to attenuate this adhesion process. Interestingly, this effect is independent of the target irradiated (that is, whether the immune cells or the endothelial cells are irradiated) [17, 21, 24].

The release of anti-adhesive cytokines also seems to be a starting point for the immunomodulatory or anti-inflammatory effect of low-dose radiotherapy [17]. Apoptosis of inflammatory cells (especially granulocytes and monocytes) likewise appears to play a role [6].

Another important target that is related to the analgesic and anti-inflammatory effects of low-dose radiotherapy is the attenuation of nitric oxide synthase and thus nitric oxide production. Nitric oxides are an important carrier of the inflammatory response and are strongly involved in inflammatory pain [17, 21].

Similar to nitric oxide synthesis, the formation of reactive oxygen metabolites, important contributors to inflammation, is also reduced [17].

What is common to these anti-inflammatory mechanisms is that their maximum effect is in the range of low-dose radiotherapy (doses between 0.3 and 1.0 Gy). At higher radiation doses, there does not seem to be an increased anti-inflammatory effect, but rather, a weaker effect can be assumed [5, 21].

These mechanisms have now been relatively well-studied at the molecular level or in cell experiments. The effect has also been demonstrated in animal models. Moreover, there are also clinical studies demonstrating that these models are applicable to patients and that low-dose radiotherapy has immunomodulatory or anti-inflammatory effects in various disease conditions [3, 5, 7, 8, 27].

### Principles of radiation physics

What radiotherapy of various degenerative and degenerative-inflammatory diseases has in common is that low radiation doses are used, usually individual doses between 0.5 and 1.0 Gy. Total doses of 3.0–6.0 Gy are usually applied. In most cases, irradiation is not performed daily, but 2–3 times a week [20, 21].

In contrast, in curative cases of malignant tumors, radiation doses of around 50 Gy are generally used in multimodal therapy approaches, and over 60 Gy in definitive radiotherapy.

Irradiation of non-malignant diseases can be performed with a linear accelerator (using photon or electron radiation) or, in part, with an orthovoltage device. A linear accelerator should be used for irradiation of activated TMJ osteoarthritis. As part of

the planning, 3D CT planning should be integrated – ideally MRI-guided – in order to protect surrounding tissue and organs at risk [2,21].

### Current evidence on radiotherapy for osteoarthritis


Since the 1920s, there have been a large number of publications showing the benefit of radiotherapy in patients with osteoarthritis. As of 2019, the author's research group has identified a total of 50 publications with over 12,000 patients who have been evaluated [11, 15]. In recent years, however, there have been 2 randomized studies on knee osteoarthritis and finger polyarthrosis, each with a smaller number of patients, that failed to show a significant benefit [13, 14]. This is in contrast to the many other studies that point to a benefit for patients – all the more so because there are also several studies that have used objective criteria to demonstrate responsiveness [8, 19]. In this regard, one assumption as to why these two randomized trials failed to show a significant benefit is that they probably included many patients with very advanced osteoarthritis. It is likely that several different pain components play a role in osteoarthritis. In addition to pain that is purely degenerative, there is also an important inflammatory pain component in activated osteoarthritis, respectively inflammatory or degenerative-inflammatory arthropathy. It can be assumed that radiotherapy can help improve the inflammatory pain component, in particular, but its effect on the purely degenerative pain component is insufficient [8, 19, 21, 27].

### Clinical effect

Many studies have shown that the analgesic, respectively the anti-inflammatory, effect occurs with a certain latency [11, 18, 28], usually ranging between 2 and 6 weeks after the end of radiotherapy. Only a small proportion of patients experience an improvement in pain during or immediately after radiotherapy. The pain minimum is usually reached between 4 and 6 weeks after the end of radiotherapy. The evaluation of the success is therefore only possible with a time delay.



**Figure 2** Design of a modern linear accelerator that can be used for the irradiation of malignant and non-malignant diseases.



**Figure 3** Positioning of the patient with a custom-made thermoplastic mask. The mask is used to immobilize the patient. A laser-guided coordinating system (drawn on the mask) helps align the patient correctly for irradiation.

### Risks of radiotherapy in the treatment of degenerative and degenerative-inflammatory diseases

Since low-dose radiotherapy is used for treatment – as mentioned above – so-called deterministic toxicities are not to be expected. These include typical side effects such as radiodermatitis or mucositis. The typical late side effects of higher-dose radiotherapy such as fibrosis or xerostomia are also not to be expected, or can be virtually ruled out, as the tolerance doses of the organs at risk are generally not reached or by far not reached [17, 21, 23].

In principle, there is a risk of stochastic side effects (e.g. tumor induction) with each application of radiation on humans. This risk of tumor induction is low in the case of radiotherapy for degenerative and degenerative-inflammatory diseases. The effective dose model can be used to estimate this risk. However, individual factors such as gender and age of the patient must still be accounted for in the assessment. The potential chance of improvement and the risks of alternative therapy options must be weighed against this risk [2, 21, 26].

For example, there is a significant risk of adverse side effects and also severe adverse side effects when analgesic drugs are taken for a longer period of time. In many cases, this risk outweighs the risk of radiotherapy [1].

In a calculation according to ICRP 2008, for an intermediate-aged patient who is irradiated unilaterally at the TMJ with a dose of  $6 \times 0.5$  Gray, there is an estimated lifetime risk of approximately 0.008 % for the induction of a malignant tumor [2, 26].

### Practical approach

Patients with activated osteoarthritis of the TMJ may also be referred for radiotherapy by dentists [12]. It should be noted that a referral is required for radiotherapy appointments. It is also advisable to send the most important documents and information in advance (provided the patient agrees to this after being informed in accordance with data protection regulations). In particular, the previous course of the disease, the course of treatment, and available cross-sectional imaging (e.g. CT or

The first clinical follow-up examination should generally take place 10–12 weeks after the end of radiotherapy. In case of an insufficient effect or lack of response, a new course of irradiation can be performed. Clinical studies have shown the benefit of such re-irradiation in degenerative or degenerative-inflammatory diseases, even in patients who did not show a response in the first course [9, 10].

Unlike enthesopathies (tendon attachment diseases) or tendinitides (tendon inflammations), in osteoarthritis, the analgesic effect of radiotherapy is often not permanent, but only present for a few years. This is most likely explained by the degenerative pain com-

ponent of osteoarthritis. In addition, osteoarthritis constitutes a progressive clinical pattern with further progressive degeneration [11, 16, 18, 20].

Since recent data suggests that predominantly or exclusively the inflammatory pain component can be favorably influenced by low-dose radiotherapy, patients with activated osteoarthritis – ideally patients in the activated relapse of osteoarthritis – should be treated particularly [19].

Whether analgesic or anti-inflammatory radiotherapy can slow down progression has not yet been clearly proven. There are – predominantly preclinical – studies that may suggest this [3, 5, 27].

Fig. 2 +and 3: UKR

MRI) with corresponding radiological findings are very relevant.

## Conclusion

Low-dose radiotherapy for treating osteoarthritis of the TMJ is a conservative treatment option. In particular, it can be applied in cases of activated osteoarthritis as well as inflammatory or degenerative-inflammatory arthropathies. Even if patients having TMJ osteoarthritis are underrepresented in the studies to date, treatment can be provided in analogy to osteoarthritis at other sites, with the aim of achieving analgesic and anti-inflammatory effects. This is also in line with the recommendation of the S2k-Guideline of the DEGRO.

The treatment utilizes low doses of irradiation. Typical deterministic toxicities are not expected and the risk of stochastic side effects appears to be low.

## Conflict of interest

The author declares that there is no conflict of interest as defined by the guidelines of the International Committee of Medical Journal Editors.

## References

- Aygün D, Kaplan S, Odaci E, Onger ME, Altunkaynak ME: Toxicity of non-steroidal anti-inflammatory drugs: a review of melatonin and diclofenac sodium association. *Histol Histopathol* 2012; 27: 417–436
- Behr M, Fanghänel J (Hrsg): *Kranio-mandibuläre Dysfunktionen. Antworten auf Fragen aus der Praxis.* Thieme, Stuttgart 2019
- Budras KD, Hartung K, Münzer BM: Licht- und elektronenmikroskopische Untersuchungen über den Einfluss von Röntgenbestrahlung auf das Stratum synoviale des entzündeten Kniegelenks. *Berl Munch Tierarztl Wochenschr* 1986; 99: 148–152
- Cocchi U: Erfolge und Mißerfolge bei Röntgenbestrahlung nichtkrebsiger Leiden. *Strahlenther* 1943; 73: 255–305
- Frey B, Gaipal US, Sarter K et al.: Whole body low dose irradiation improves the course of beginning polyarthritis in human TNF-transgenic mice. *Autoimmunity* 2009; 42: 346–348
- Gaipal US, Meister S, Lödermann B et al.: Activation-induced cell death and total Akt content of granulocytes show a biphasic course after low-dose radiation. *Autoimmunity* 2009; 42: 340–342
- Hautmann AH, Wolff D, Hilgendorf I et al.: Total nodal irradiation in patients with severe treatment-refractory chronic graft-versus-host disease after allogeneic stem cell transplantation: Response rates and immunomodulatory effects. *Radiother Oncol* 2015; 116: 287–293
- Hautmann MG, Jung E-M, Beyer LP et al.: Ist die Strahlentherapie eine effektive Behandlungsoption der Bakerzyste? *Strahlenther Onkol* 2019; 195: 69–76
- Hautmann MG, Neumaier U, Kölbl O: Re-irradiation for painful heel spur syndrome. Retrospective analysis of 101 heels. *Strahlenther Onkol* 2014; 190: 298–303
- Hautmann MG, Rechner P, Hipp M et al.: Rebestrahlung bei Arthrose – retrospektive Analyse von 217 Gelenken. *Strahlenther Onkol* 2019; 195: 1060–1067
- Hautmann MG, Rechner P, Neumaier U et al.: Radiotherapy for osteoarthritis – an analysis of 295 joints treated with a linear accelerator. *Strahlenther Onkol* 2020; 196: 715–724
- Kassenzahnärztliche Bundesvereinigung, Spitzenverband Bund der Krankenkassen: Bundesmantelvertrag – Zahnärzte (BMV-Z) 2021
- Mahler EAM, Minten MJJ, Leseman-Hoogenboom MM et al.: Effectiveness of low-dose radiation therapy on symptoms in patients with knee osteoarthritis: a randomised, double-blinded, sham-controlled trial. *Ann Rheum Dis* 2019; 78: 83–90
- Minten MJM, Leseman-Hoogenboom MM, Kloppenburg M et al.: Lack of beneficial effects of low-dose radiation therapy on hand osteoarthritis symptoms and inflammation: a randomised, blinded, sham-controlled trial. *Osteoarthritis Cartilage* 2018; 26: 1283–1290
- Minten MJM, Mahler E, Broeder AA den, Leer JWH, van den Ende CH: The efficacy and safety of low-dose radiotherapy on pain and functioning in patients with osteoarthritis: a systematic review. *Rheumatol Int* 2016; 36: 133–142
- Mücke R, Seegenschmiedt MH, Heyd R et al.: Strahlentherapie bei schmerzhafter Kniegelenkarthrose (Gonarthrose): Ergebnisse einer deutschen Patterns-of-Care-Studie. *Strahlenther Onkol* 2010; 186: 7–17
- Mücke R, Micke O, Seegenschmiedt MH: Strahlentherapie gutartiger Erkrankungen. S2e-Leitlinie der Deutschen Gesellschaft für Radioonkologie (DEGRO). 2018
- Niewald M, Seegenschmiedt MH, Micke O et al.: Randomized, multicenter trial on the effect of radiation therapy on plantar fasciitis (painful heel spur) comparing a standard dose with a very low dose: mature results after 12 months' follow-up. *Int J Radiat Oncol Biol Phys* 2012; 84: e455–62
- Ott OJ, Micke O, Mücke R et al.: Low-dose radiotherapy: Mayday, mayday. We've been hit! *Strahlenther Onkol* 2019; 195: 285–288
- Ott OJ, Niewald M, Weitmann H-D et al.: DEGRO guidelines for the radiotherapy of non-malignant disorders. Part II: Painful degenerative skeletal disorders. *Strahlenther Onkol* 2015; 191: 1–6
- Reichl B, Block A, Schäfer U et al.: DEGRO practical guidelines for radiotherapy of non-malignant disorders: Part I: Physical principles, radiobiological mechanisms, and radiogenic risk. *Strahlenther Onkol* 2015; 191: 701–709
- Rödel F, Hofmann D, Auer J et al.: The anti-inflammatory effect of low-dose radiation therapy involves a diminished CCL20 chemokine expression and granulocyte/endothelial cell adhesion. *Strahlenther Onkol* 2008; 184: 41–47
- Ruppert R, Seegenschmiedt MH, Sauer R: Radiotherapie von Arthrosen. Indikation, Technik, klinische Ergebnisse. *Orthopade* 2004; 33: 56–62
- Seegenschmiedt MH, Makoski H-B, Trott K-R et al.: Radiotherapy for non-malignant disorders. Springer, Berlin, Heidelberg 2008
- Seegenschmiedt MH, Micke O, Willich N: Radiation therapy for nonmalignant diseases in Germany. Current concepts and future perspectives. *Strahlenther Onkol* 2004; 180: 718–730
- The 2007 Recommendations of the International Commission on Radiological Protection. ICRP publication 103. *Ann ICRP* 2007; 37: 1–332
- Trott KR, Parker R, Seed MP: Die Wirkung von Röntgenstrahlen auf die experimentelle Arthritis der Ratte. *Strahlenther Onkol* 1995; 171: 534–538
- Zschache H: Ergebnisse der Röntgen-schwachbestrahlung. *Radiol Diagn (Berl)* 1972; 13: 181–186



Photo: UKR

**PD DR. MATTHIAS G. HAUTMANN,**  
Clinic and Polyclinic for Radiotherapy,  
University Hospital Regensburg,  
93042 Regensburg